

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
6 October 2005 (06.10.2005)

PCT

(10) International Publication Number
WO 2005/093101 A1

(51) International Patent Classification⁷: C12Q 1/68, 1/68

(21) International Application Number:

PCT/IB2005/000771

(22) International Filing Date: 24 March 2005 (24.03.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

0406863.1

26 March 2004 (26.03.2004)

GB

(71) Applicant (for all designated States except US): **QIAGEN AS** [NO/NO]; Frysjavaeien 40, N-0884 Oslo (NO).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **LARSEN, Frank** [NO/NO]; Gaustadveien 93, N-0372 Oslo (NO).

(74) Common Representative: **QIAGEN AS**; c/o Page White & Farrer, 54 Doughty Street, London WC1N 2LS (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

[Continued on next page]

(54) Title: NUCLEIC ACID SEQUENCING



(57) Abstract: A method for determining a target nucleic acid sequence, wherein the target nucleic acid sequence is comprised in a preparation comprising a non-target nucleic acid sequence, the target nucleic acid sequence and the non-target nucleic acid sequence each having a first region of common sequence upstream of a first region of dissimilar sequence upstream of a second region of dissimilar sequence, the method comprising: (a) contacting the preparation with a blocking oligonucleotide complementary to at least a portion of the first region of dissimilar sequence of the non-target nucleic acid sequence, under conditions to hybridise the blocking oligonucleotide thereto; (b) contacting the preparation with a sequencing primer complementary to at least a portion of the first region of common sequence, under conditions to hybridise the primer to the target nucleic acid sequence; and (c) subjecting the preparation to a sequencing reaction, such that the sequencing reaction proceeds into the second region of dissimilar sequence of the target nucleic acid sequence, thereby determining at least the second region of dissimilar sequence of the target nucleic acid sequence; and wherein the blocking oligonucleotide blocks the sequencing reaction at least from proceeding into the second region of dissimilar sequence of the non-target nucleic acid sequence.

WO 2005/093101 A1



— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.